

NSW Threatened Species Scientific Committee

Notice of Preliminary Determination

The NSW Threatened Species Scientific Committee, established under the *Biodiversity Conservation Act 2016* (the Act), has made a Preliminary Determination to support a proposal to remove *Xanthosia scopulicola* J.M.Hart & Henwood from the Schedules of the Act by omitting reference to this species from Part 3 of Schedule 1 (Vulnerable species). The omission of species from the Schedules is provided for by Part 4 of the Act.

How to make a submission

The NSW TSSC welcomes public involvement in the assessment process and places preliminary determinations on public exhibition on the NSW TSSC pages on the NSW Department of Climate Change, Energy, the Environment and Water (NSW DCCEEW) website. This public exhibition provides an opportunity for the public to comment on this preliminary determination as well as provide any additional information that is relevant to the assessment.

Postal submissions regarding this Preliminary Determination may be sent to:

Secretariat
NSW Threatened Species Scientific Committee
Locked Bag 5022
Parramatta NSW 2124.

Email submissions in Microsoft Word or PDF formats to:

scientific.committee@environment.nsw.gov.au

Submissions close 13 December 2024

What happens next?

After considering any submissions received during the public exhibition period the NSW TSSC will make a Final Determination and a notice will be placed on the NSW DCCEEW website to announce the outcome of the assessment. If the Final Determination is to support a listing, then it will be added to the Schedules of the Act when the Final Determination is published on the legislation website. www.legislation.nsw.gov.au.

Privacy information

The information you provide in your submission may be used by the NSW TSSC in the assessment to determine the conservation status and listing or delisting of threatened or extinct species, threatened populations and threatened or collapsed ecological communities or to assess key threatening processes.

The NSW TSSC may be asked to share information on assessments with NSW Government agencies, the Commonwealth Government and other State and Territory governments to collaborate on national threatened species assessments using a common assessment method and to assist in the management of species and ecological communities.

NSW Threatened Species Scientific Committee

If your submission contains information relevant to the assessment it may be provided to state and territory government agencies and scientific committees as part of this collaboration.

If you wish your identity and personal information in your submission to be treated as confidential you must:

- *request your name be treated as confidential, and*
- *not include any of your personal information in the main text of the submission or attachments so that it can be easily removed.*

Senior Professor Kristine French
Chairperson
NSW Threatened Species Scientific Committee

Public Exhibition period: 13/09/2024 - 13/12/2024

Preliminary Determination

The NSW Threatened Species Scientific Committee, established under the *Biodiversity Conservation Act 2016* (the Act), has made a Preliminary Determination to support a proposal to remove *Xanthosia scopulicola* J.M.Hart & Henwood from the Schedules of the Act by omitting reference to this species from Part 3 of Schedule 1 (Vulnerable species). The omission of species from the Schedules is provided for by Part 4 of the Act.

Summary of Conservation Assessment

The NSW Threatened Species Scientific Committee has found that:

1. *Xanthosia scopulicola* J.M.Hart & Henwood (family Apiaceae) was described by Hart and Henwood (2000) as an “ascending perennial subshrub to 20 cm high. Taproot woody with a flaky or corky surface. Plants stellately tomentose on young stems and leaves; the mature stems exorticating. Leaves grey-green, cauline, juvenile leaves trifoliolate, adult leaves simple. Petiole 2–15 mm long, angular, grooved above, sheathing shortly at the base, sheaths 1–2 mm long. Leaflets of the trifoliolate leaves ovate, the terminal leaflet longer than the lateral leaflets, terminal leaflet 5–20 mm long, 5–25 mm wide, lateral leaflets 3–5 mm long, 3–4 mm wide, petiolulate or sessile. Simple leaves ovate, 5–20 mm long, 5–25 mm wide. Leaf margins flat to slightly recurved, crenate; apex obtuse. Inflorescence a compound umbel with 2 or 3 rays, 1–4 flowers per ray; flowers bisexual or rarely male. Rays terete, 2–4 mm long. Involucral bracts 2 or 3, triangular, foliaceous, green, longer or shorter than the rays, 2–3 mm long, c. 1 mm wide, apex acute. Bracteoles 3, elliptic to obovate, petaloid, yellow, shorter or longer than flowers, 3–5 mm long, 1–3 mm wide, apex acuminate or obtuse. Inflorescences pedunculate; peduncles 5–25 mm long. Flowers pedicellate; pedicels 1–2 mm long. Sepals ovate, 1–1.2 mm long, c. 0.7 mm wide, green or yellow, base truncate, apex obtuse, glabrous. Petals longer than sepals, 1.0–1.2 mm long, c. 0.5 mm wide, white, midrib adaxially keeled and bridged with the inflexed appendage, apex obtuse, appendage smooth. Nectaries raised and prominent, c. 0.9 mm high, white or yellow, hirsute. Styles 1.3–2.3 mm long. Ovary hirsute. Male flowers having an undeveloped inconspicuous ovary, with the styles barely protruding above the nectaries. Fruit brown, ovate, 2.0–2.2 mm long, 2.0–2.2 mm wide, c. 0.5 mm deep. Monocarps hirsute on the summit, elliptic in transverse section, 7–9-ribbed; ribs keeled.”
2. *Xanthosia scopulicola* is restricted to sandstone cliff faces or rocky outcrops of the upper Blue Mountains. It is found along approximately 25 km of escarpment in the Jamison and Kedumba Valleys between Kings Tableland in Wentworth Falls and the Megalong Cleft in Katoomba, and on the northern side of the Grose Valley. In 2019, the species was recorded for the first time from the northern escarpment of the Grose Valley, where it has now been found at three sites with a linear extent of 10 km. Most of the population is located within Blue Mountains National Park.
3. The geographic distribution of *Xanthosia scopulicola* is highly restricted. The Extent of Occurrence (EOO) is estimated at 183 km², and the area of occurrence (AOO) is estimated at 52 km². The Extent of Occurrence (EOO) is based on a

NSW Threatened Species Scientific Committee

minimum convex polygon enclosing all mapped occurrences of the species, the method of assessment recommended by IUCN (2024). The AOO is based on 2 x 2 km grid cells, the scale recommended for assessing area of occupancy by IUCN (2024).

4. The total population size of *Xanthosia scopulicola* is estimated to be a minimum of 2,000. The population estimate is based on an extrapolation of counts from surveys along walking tracks targeting known records (V. Wong pers. obs. October 2022). Based on the potential for gene exchange between individuals, there are considered to be two populations of *Xanthosia scopulicola*: one in the Grose Valley, and one in Jamieson/Kedumba Valleys.
5. *Xanthosia scopulicola* grows in skeletal soils in cracks and crevices of sandstone cliff faces, on ledges, under overhangs or in clay bands in the rockface, or (less commonly) on rocky outcrops above the cliffs (ALA 2024; National Herbarium of New South Wales collections database accessed 20 February 2024). Known occurrences are predominantly on south-facing cliffs along the northern escarpments of the Grose and Jamieson Valleys in the Blue Mountains NP, at altitudes of (550–)750–1000 m elevation. Where aspect is included in occurrence records, south facing is the most commonly given, although aspect varies, presumably due to undulations in the cliff face. Mean annual rainfall at Katoomba (the nearest weather station, 1017 m a.s.l.) is 1410 mm (Bureau of Meteorology 2024). Associated species include *Alania cunninghamii*, *Amperea xiphoclada*, *Baeckea linifolia*, *Blandfordia cunninghamii*, *Dracophyllum secundum*, *Empodisma minus*, *Epacris rigida*, *Eucalyptus oreades* and *E. piperita*, *Gleichenia dicarpa*, *G. rupestris*, and *Sprengelia monticola* (ALA 2024; National Herbarium of New South Wales collections database accessed 20 February 2024).
6. *Xanthosia scopulicola* is a perennial subshrub with a deep taproot and a compact and much-branched habit (Hart and Henwood 2000; PlantNET 2024). There is no information available on longevity or time to maturity. However, the slow development of the exfoliating bark over at least several years (V. Wong pers. obs. July 2023) suggests that the subshrub can live for many years. Pollination mechanisms of eastern Australian *Xanthosia* are unknown but, as the flowers of *X. scopulicola* are inconspicuous it is inferred they may be largely self-pollinated. The fruit dehisces into two single-seeded indehiscent mericarps, which are the unit of dispersal.
7. *Xanthosia scopulicola* at Narrow Neck was observed to resprout after fire and to have very limited recruitment from seed post-fire, contrasting with the co-occurring *X. pilosa*, which was observed to have mass recruitment from seed after the same fire in late 2019 (V. Wong pers. obs. December 2021). Resprouting individuals have been observed to have flowers two years after fire but are likely to flower in the first year after fire (V. Wong pers. obs. December 2021).
8. All known individuals of *Xanthosia scopulicola* occur within two expanses of habitat, along the (predominantly northern) escarpments of the Jamieson and Kedumba Valleys and along the northern escarpment of the Grose Valley. These two areas constitute two threat-defined locations, based on the most serious plausible threat of adverse fire regimes. The Jamieson/Kedumba Valleys and the Grose Valley are not only geographically separated, their different fire histories also indicate they are unlikely to be affected by the same fire event.

NSW Threatened Species Scientific Committee

9. Previously documented threats to *Xanthosia scopulicola* include habitat disturbance from track maintenance, disturbance by walkers, urban runoff and weed encroachment. Most of these threats are associated with walking tracks and urban drain outflows within known habitat, which is strongly suspected to be a very small proportion of the total extent of this species. Other plausible threats to *Xanthosia scopulicola* include adverse fire regimes, and increased frequency and intensity of both drought and extreme rainfall events due to climate change. However, the ecology of the species, especially its preference for open cliff-face habitats, and evidence from field observations undertaken for this assessment and other available data, mean that continuing decline is not considered to be occurring due to any of these potential threats, and there is not a high degree of certainty that any future declines will occur.
10. Continuing decline is not evident in the known populations of *Xanthosia scopulicola*. Due to the preferred habitat of largely unvegetated, high, south-facing cliffs that the species occupies, neither high frequency or high severity fire are considered to be causing declines, nor are they able to be considered likely to do so in the future. The species ecology and habitat preference also mean the species is resilient to the effects of drought and extreme rainfall events, and no evidence of these mechanisms causing non-trivial declines is known. All other threats, including anthropogenic disturbance, urban runoff and weed invasion are considered to be trivial in nature. Field observations carried out for this assessment and other available data do not demonstrate an observed, estimated, inferred, or projected continuing decline in EOO, AOO, area, extent and/or quality of habitat, number of locations or populations, or number of mature individuals for *X. scopulicola*, and there is little confidence that such declines may become apparent in the future.
11. The identified plausible threats to *Xanthosia scopulicola* are likely to be localised in nature if they do become apparent in the future. Thus, these threats are not considered likely to rapidly drive the species to extinction in a very short time (1-2 generations) across its full distribution.
12. In view of the above, the NSW Threatened Species Scientific Committee is of the opinion that *Xanthosia scopulicola* J.M.Hart & Henwood is not eligible to be listed as a threatened species in any category under the Act.

Assessment against *Biodiversity Conservation Regulation 2017* criteria

The Clauses used for assessment are listed below for reference.

Overall Assessment Outcome: *Xanthosia scopulicola* J.M.Hart & Henwood was found to be ineligible for listing as a threatened species as none of the criteria were met.

Clause 4.2 – Reduction in population size of species (Equivalent to IUCN criterion A)

Assessment Outcome: Data Deficient

(1) - The species has undergone or is likely to undergo within a time frame appropriate to the life cycle and habitat characteristics of the taxon:

	(a)	for critically endangered species	a very large reduction in population size, or
--	-----	-----------------------------------	-----------------------------------------------

NSW Threatened Species Scientific Committee

	(b)	for endangered species	a large reduction in population size, or
	(c)	for vulnerable species	a moderate reduction in population size.
(2) - The determination of that criteria is to be based on any of the following:			
	(a)	direct observation,	
	(b)	an index of abundance appropriate to the taxon,	
	(c)	a decline in the geographic distribution or habitat quality,	
	(d)	the actual or potential levels of exploitation of the species,	
	(e)	the effects of introduced taxa, hybridisation, pathogens, pollutants, competitors or parasites.	

Clause 4.3 – Restricted geographic distribution of species and other conditions (Equivalent to IUCN criterion B)

Assessment Outcome: Clause not met

The geographic distribution of the species is:			
	(a)	for critically endangered species	very highly restricted, or
	(b)	for endangered species	highly restricted, or
	(c)	for vulnerable species	moderately restricted.
and at least 2 of the following 3 conditions apply:			
	(d)	the population or habitat of the species is severely fragmented or nearly all the mature individuals of the species occur within a small number of locations,	
	(e)	there is a projected or continuing decline in any of the following:	
		(i)	an index of abundance appropriate to the taxon,
		(ii)	the geographic distribution of the species,
		(iii)	habitat area, extent or quality,
		(iv)	the number of locations in which the species occurs or of populations of the species.
	(f)	extreme fluctuations occur in any of the following:	
		(i)	an index of abundance appropriate to the taxon,
		(ii)	the geographic distribution of the species,
		(iii)	the number of locations in which the species occur or of populations of the species.

Clause 4.4 – Low numbers of mature individuals of species and other conditions

(Equivalent to IUCN criterion Clause C)

Assessment Outcome: Clause not met

The estimated total number of mature individuals of the species is:			
	(a)	for critically endangered species	very low, or
	(b)	for endangered species	low, or
	(c)	for vulnerable species	moderately low.
and either of the following 2 conditions apply:			
	(d)	a continuing decline in the number of mature individuals that is (according to an index of abundance appropriate to the species):	
		(i)	for critically endangered species very large, or

NSW Threatened Species Scientific Committee

	(ii)	for endangered species	large, or
	(iii)	for vulnerable species	moderate,
(e)	both of the following apply:		
	(i)	a continuing decline in the number of mature individuals (according to an index of abundance appropriate to the species), and	
	(ii)	at least one of the following applies:	
	(A)	the number of individuals in each population of the species is:	
		(I)	for critically endangered species extremely low, or
		(II)	for endangered species very low, or
		(III)	for vulnerable species low,
	(B)	all or nearly all mature individuals of the species occur within one population,	
	(C)	extreme fluctuations occur in an index of abundance appropriate to the species.	

**Clause 4.5 – Low total numbers of mature individuals of species
(Equivalent to IUCN criterion D)**

Assessment Outcome: Clause not met

The total number of mature individuals of the species is:			
	(a)	for critically endangered species	extremely low, or
	(b)	for endangered species	very low, or
	(c)	for vulnerable species	low.

**Clause 4.6 – Quantitative analysis of extinction probability
(Equivalent to IUCN criterion E)**

Assessment Outcome: Data Deficient

The probability of extinction of the species is estimated to be:			
	(a)	for critically endangered species	extremely high, or
	(b)	for endangered species	very high, or
	(c)	for vulnerable species	high.

**Clause 4.7 – Very highly restricted geographic distribution of species–
vulnerable species**

(Equivalent to IUCN criterion D2)

Assessment Outcome: Clause not met

For vulnerable species,	the geographic distribution of the species or the number of locations of the species is very highly restricted such that the species is prone to the effects of human activities or stochastic events within a very short time period.
-------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Senior Professor Kristine French
Chairperson
NSW Threatened Species Scientific Committee

NSW Threatened Species Scientific Committee

Supporting Documentation:

Wong, V, Gibbons K.L. (2024) Conservation Assessment of *Xanthosia scopulicola* J.M.Hart & Henwood (Apiaceae). NSW Threatened Species Scientific Committee.

References:

Atlas of Living Australia (ALA) (2024) *Xanthosia scopulicola* occurrence records [dataset]. Available at:

<https://bie.ala.org.au/species/https://id.biodiversity.org.au/node/apni/2898254> (accessed 20 February 2024).

Bureau of Meteorology (2024) Climate statistics for Australian locations. Available at: http://www.bom.gov.au/climate/averages/tables/cw_063039.shtml (accessed 13 February 2024).

Hart JM, Henwood MJ (2000) Systematics of the *Xanthosia pilosa* complex (Apiaceae: Hydrocotyloideae). *Australian Systematic Botany* **13(2)**,245–66.

IUCN Standards and Petitions Subcommittee (2024) Guidelines for Using the IUCN Red List Categories and Criteria Version 16 (March 2024). Available at: <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> (accessed on 18 April 2024).

PlantNET 2024 (The NSW Plant Information Network System). Royal Botanic Gardens and Domain Trust, Sydney. <https://plantnet.rbgsyd.nsw.gov.au> (accessed 20 February 2024).