

Ecosystem credit demand estimate


Offset Trading Group indicative longer-term demand by IBRA region and subregion

The following ecosystem credit demand estimate has been produced using a spatial model and publicly available planning information. This is presented as demand per Offset Trading Group by IBRA (Interim Biogeographic Regionalisation for Australia) region and subregion. The estimate is based on information as of 28 March 2023. Landholders should seek their own legal, financial and ecological advice before participating in the market. As the model ingests data from very early in the planning processes for major projects, before a biodiversity development assessment report has been undertaken, the development footprints and resulting credit predictions associated with early-stage projects can be highly uncertain.

The data are presented in a series of tables, one for each IBRA bioregion as follows (click on the bioregion of interest to jump to the table):

Table 1	Brigalow Belt South Bioregion	2
Table 2	Broken Hill Complex Bioregion	6
Table 3	Channel Country Bioregion	7
Table 4	Cobar Peneplain Bioregion	8
Table 5	Darling Riverine Plains Bioregion	9
Table 6	Mulga Lands Bioregion	11
Table 7	Murray Darling Depression Bioregion	11
Table 8	Nandewar Bioregion	12
Table 9	New England Tablelands Bioregion	14
Table 10	NSW North Coast Bioregion	18
Table 11	NSW South Western Slopes Bioregion	23
Table 12	Riverina Bioregion	26
Table 13	Simpson Strzelecki Dunefields Bioregion	28
Table 14	South East Corner Bioregion	29
Table 15	South Eastern Highlands Bioregion	30
Table 16	South Eastern Queensland Bioregion	36
Table 17	Sydney Basin Bioregion	38

A numerical credit estimate is provided as an average, with a range presented in parentheses (). The following categories have also been used to depict demand:



An Excel spreadsheet containing the same data is also available for download.

Table 1 Brigalow Belt South Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Liverpool Plains		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Brigalow within the Brigalow Belt South, Nandewar and Darling Riverine Plains Bioregions	Not detected	0
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Not detected	0
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Moderate	1,286 (631 to 1,941)
Inland Riverine Forests greater than or equal to 50% and less than 70%	Low	12 (7 to 17)
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Northern Montane Heaths less than 50%	Low	5 (5 to 6)
North-west Alluvial Sand Woodlands greater than or equal to 50% and less than 70%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Low	34 (20 to 49)
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Low	115 (66 to 163)
Pilliga Outwash Dry Sclerophyll Forests less than 50%	Low	571 (367 to 774)
Semi-evergreen Vine Thicket in the Brigalow Belt South and Nandewar Bioregions	Low	31 (18 to 43)
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Moderate	2,028 (1,396 to 2,660)
Western Slopes Dry Sclerophyll Forests less than 50%	Moderate	1,212 (834 to 1,590)
Western Slopes Grassy Woodlands greater than or equal to 70% and less than 90%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North	Low	830 (481 to 1,178)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions		
Liverpool Range		
Coastal Valley Grassy Woodlands less than 50%	Low	38 (20 to 56)
Eastern Riverine Forests less than 50%	Not detected	0
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	115 (79 to 151)
Western Slopes Dry Sclerophyll Forests less than 50%	Low	<1
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	994 (486 to 1,502)
Narrandool		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Northern Basalts		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Brigalow within the Brigalow Belt South, Nandewar and Darling Riverine Plains Bioregions	Not detected	0
Carbeen Open Forest Community in the Darling Riverine Plains and Brigalow Belt South Bioregions	Not detected	0
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Not detected	0
Native Vegetation on Cracking Clay Soils of the Liverpool Plains	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Pilliga Outwash Dry Sclerophyll Forests less than 50%	Not detected	0
Semi-evergreen Vine Thicket in the Brigalow Belt South and Nandewar Bioregions	Not detected	0
Subtropical Semi-arid Woodlands less than 50%	Not detected	0
Western Peneplain Woodlands less than 50%	Not detected	0
Western Slopes Dry Sclerophyll Forests less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Northern Outwash		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Brigalow within the Brigalow Belt South, Nandewar and Darling Riverine Plains Bioregions	Not detected	0
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Not detected	0
Pilliga Outwash Dry Sclerophyll Forests less than 50%	Not detected	0
Western Slopes Dry Sclerophyll Forests less than 50%	Not detected	0
Pilliga		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Low	8 (5 to 12)
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Not detected	0
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
Pilliga Outwash Dry Sclerophyll Forests less than 50%	Not detected	0
Riverine Plain Grasslands greater than or equal to 50% and less than 70%	Low	3 (3 to 4)
Western Slopes Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Not detected	0
Western Slopes Dry Sclerophyll Forests less than 50%	Strong	6,192 (4,263 to 8,121)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	871 (599 to 1,142)
Pilliga Outwash		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Brigalow within the Brigalow Belt South, Nandewar and Darling Riverine Plains Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Not detected	0
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Low	46 (23 to 70)
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Not detected	0
Inland Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Not detected	0
North-west Alluvial Sand Woodlands greater than or equal to 50% and less than 70%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Pilliga Outwash Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Pilliga Outwash Dry Sclerophyll Forests less than 50%	Low	513 (330 to 695)
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Western Slopes Dry Sclerophyll Forests less than 50%	Strong	11,976 (8,245 to 15,707)
Western Slopes Grassy Woodlands greater than or equal to 70% and less than 90%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Talbragar Valley		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Low	299 (146 to 452)
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Moderate	2,713 (1,537 to 3,890)
Inland Riverine Forests greater than or equal to 50% and less than 70%	Low	151 (86 to 217)
Inland Rocky Hill Woodlands less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Low	20 (12 to 29)
Western Slopes Dry Sclerophyll Forests less than 50%	Moderate	1,181 (813 to 1,549)
Western Slopes Grassy Woodlands greater than or equal to 50% and less than 70%	Low	73 (36 to 110)
Western Slopes Grassy Woodlands less than 50%	Strong	9,729 (4,757 to 14,703)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Moderate	1,975 (1,010 to 2,940)

Table 2 Broken Hill Complex Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Barrier Range		
Acacia loderi Shrublands	Strong	8,631 (5,192 to 12,071)
Aeolian Chenopod Shrublands less than 50%	Strong	36,914 (23,755 to 50,067)
Gibber Chenopod Shrublands greater than or equal to 50% and less than 70%	Strong	72,415 (32,403 to 112,412)
Gibber Chenopod Shrublands less than 50%	Strong	47,300 (21,165 to 73,424)
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Low	67 (41 to 93)
Inland Saline Lakes less than 50%	Low	37 (23 to 50)
North-west Floodplain Woodlands less than 50%	Low	127 (78 to 177)
Porcupine Grass—Red Mallee—Gum Coolabah hummock grassland/low sparse woodland in the Broken Hill Complex Bioregion	Low	287 (150 to 424)
Riverine Chenopod Shrublands less than 50%	Strong	6,817 (6,008 to 7,624)
Sand Plain Mulga Shrublands greater than or equal to 50% and less than 70%	Strong	30,294 (18,222 to 42,366)
Semi-arid Floodplain Grasslands greater than or equal to 50% and less than 70%	Strong	11,535 (6,338 to 16,729)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Semi-arid Sand Plain Woodlands greater than or equal to 50% and less than 70%	Low	18 (8 to 28)
Stony Desert Mulga Shrublands less than 50%	Strong	11,631 (6,075 to 17,188)
Barrier Range Outwash		
Acacia loderi Shrublands	Moderate	1,561 (939 to 2,183)
Aeolian Chenopod Shrublands less than 50%	Strong	9,272 (5,967 to 12,575)
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Gibber Chenopod Shrublands greater than or equal to 50% and less than 70%	Strong	11,242 (5,030 to 17,451)
Gibber Chenopod Shrublands less than 50%	Moderate	1,025 (459 to 1,591)
Inland Floodplain Shrublands greater than or equal to 50% and less than 70%	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Inland Saline Lakes less than 50%	Not detected	0
North-west Floodplain Woodlands less than 50%	Low	75 (46 to 104)
Sand Plain Mulga Shrublands greater than or equal to 50% and less than 70%	Low	467 (281 to 653)
Sand Plain Mulga Shrublands less than 50%	Moderate	1,053 (634 to 1,473)
Semi-arid Floodplain Grasslands greater than or equal to 50% and less than 70%	Low	800 (440 to 1,160)
Semi-arid Sand Plain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Stony Desert Mulga Shrublands less than 50%	Strong	5,461 (2,852 to 8,070)

Table 3 Channel Country Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Core Ranges		
Desert Woodlands less than 50%	Low	16 (9 to 25)
Gibber Chenopod Shrublands greater than or equal to 50% and less than 70%	Low	8 (3 to 12)
Gibber Chenopod Shrublands less than 50%	Low	30 (14 to 47)
Gibber Transition Shrublands less than 50%	Low	19 (10 to 28)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Riverine Chenopod Shrublands less than 50%	Low	229 (202 to 256)
Sturt Stony Desert		
Gibber Transition Shrublands less than 50%	Low	22 (11 to 32)
North-west Floodplain Woodlands less than 50%	Low	25 (15 to 34)
Riverine Chenopod Shrublands less than 50%	Low	713 (628 to 797)
Sand Plain Mulga Shrublands greater than or equal to 50% and less than 70%	Low	59 (36 to 83)
Sand Plain Mulga Shrublands less than 50%	Low	71 (43 to 99)

Table 4 Cobar Peneplain Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Barnato Downs		
Acacia loderi Shrublands	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Stony Desert Mulga Shrublands less than 50%	Not detected	0
Western Peneplain Woodlands less than 50%	Not detected	0
Canbelego Downs		
Acacia loderi Shrublands	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Floodplain Transition Woodlands greater than or equal to 70% and less than 90%	Low	15 (9 to 22)
Inland Floodplain Swamps greater than or equal to 50% and less than 70%	Low	6 (3 to 10)
Inland Rocky Hill Woodlands less than 50%	Moderate	1,413 (884 to 1,941)
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
Sand Plain Mallee Woodlands greater than or equal to 50% and less than 70%	Low	5 (4 to 6)
Sand Plain Mallee Woodlands greater than or equal to 70% and less than 90%	Not detected	0
Western Peneplain Woodlands greater than or equal to 50% and less than 70%	Low	634 (287 to 981)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Western Peneplain Woodlands less than 50%	Moderate	1,945 (881 to 3,010)
Lachlan Plains		
Acacia loderi Shrublands	Not detected	0
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Not detected	0
Inland Riverine Forests less than 50%	Low	<1
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Western Peneplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Western Peneplain Woodlands less than 50%	Not detected	0
Nymagee		
Acacia loderi Shrublands	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Not detected	0
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
Sand Plain Mallee Woodlands greater than or equal to 70% and less than 90%	Not detected	0
Stony Desert Mulga Shrublands less than 50%	Not detected	0
Western Peneplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Western Peneplain Woodlands less than 50%	Not detected	0
Western Slopes Dry Sclerophyll Forests less than 50%	Not detected	0

Table 5 Darling Riverine Plains Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Bogan-Macquarie		

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Not detected	0
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Not detected	0
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Castlereagh-Barwon		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Floodplain Shrublands greater than or equal to 70% and less than 90%	Not detected	0
Louth Plains		
Aeolian Chenopod Shrublands less than 50%	Low	3 (2 to 4)
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Low	272 (166 to 378)
Riverine Chenopod Shrublands less than 50%	Low	115 (101 to 129)
Semi-arid Floodplain Grasslands greater than or equal to 50% and less than 70%	Low	4 (2 to 5)
Pooncarie-Darling		
Acacia loderi Shrublands	Not detected	0
Acacia melvillei Shrubland in the Riverina and Murray-Darling Depression Bioregions	Not detected	0
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Riverine Forests less than 50%	Not detected	0
Semi-arid Sand Plain Woodlands less than 50%	Not detected	0

Table 6 Mulga Lands Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Paroo Overflow		
Aeolian Chenopod Shrublands less than 50%	Low	20 (13 to 27)
Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions	Low	139 (85 to 193)
Gibber Transition Shrublands less than 50%	Low	16 (8 to 23)
Paroo-Darling Sands		
Acacia loderi Shrublands	Not detected	0
Aeolian Chenopod Shrublands less than 50%	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Gibber Chenopod Shrublands less than 50%	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Not detected	0
Inland Saline Lakes less than 50%	Not detected	0
North-west Floodplain Woodlands less than 50%	Not detected	0

Table 7 Murray Darling Depression Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Darling Depression		
Acacia loderi Shrublands	Not detected	0
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Not detected	0
Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes Bioregions	Not detected	0
Riverine Plain Grasslands less than 50%	Not detected	0
South Olary Plain		
Acacia melvillei Shrubland in the Riverina and Murray-Darling Depression Bioregions	Not detected	0
Aeolian Chenopod Shrublands less than 50%	Strong	19,162 (12,332 to 25,990)
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Dune Mallee Woodlands less than 50%	Strong	6,287 (4,597 to 7,977)
Gibber Chenopod Shrublands less than 50%	Strong	17,850 (7,987 to 27,709)
Inland Floodplain Shrublands greater than or equal to 50% and less than 70%	Not detected	0
Inland Floodplain Shrublands less than 50%	Low	218 (143 to 294)
Inland Floodplain Swamps greater than or equal to 50% and less than 70%	Low	6 (3 to 10)
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Strong	5,219 (3,183 to 7,257)
Inland Riverine Forests less than 50%	Not detected	0
Inland Saline Lakes less than 50%	Strong	11,703 (7,271 to 16,137)
Riverine Chenopod Shrublands greater than or equal to 50% and less than 70%	Strong	5,221 (4,602 to 5,839)
Riverine Chenopod Shrublands less than 50%	Low	88 (78 to 99)
Riverine Plain Grasslands less than 50%	Moderate	1,014 (832 to 1,195)
Sand Plain Mallee Woodlands greater than or equal to 50% and less than 70%	Low	586 (327 to 845)
Sand Plain Mallee Woodlands less than 50%	Strong	60,336 (45,095 to 75,576)
Sand Plain Mulga Shrublands greater than or equal to 50% and less than 70%	Moderate	2,655 (1,597 to 3,713)
Sand Plain Mulga Shrublands less than 50%	Strong	13,677 (8,227 to 19,127)
Sandhill Pine Woodland in the Riverina, Murray-Darling Depression and NSW South Western Slopes Bioregions	Moderate	4,087 (2,254 to 5,919)
Semi-arid Sand Plain Woodlands greater than or equal to 50% and less than 70%	Strong	14,356 (6,092 to 22,618)
Semi-arid Sand Plain Woodlands less than 50%	Low	753 (320 to 1,187)
Tecticornia lylei, Wiry Glasswort, low open-shrubland in the Murray Darling Depression Bioregion	Not detected	0

Table 8 Nandewar Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Nandewar Northern Complex		
Eastern Riverine Forests less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Inland Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Low	13 (7 to 18)
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 70% and less than 90%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Western Slopes Grassy Woodlands greater than or equal to 70% and less than 90%	Low	17 (8 to 25)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	1 (1 to 2)
Peel		
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Moderate	1,081 (589 to 1,573)
Mount Kaputar high elevation and dry rainforest land snail and slug community in the Nandewar and Brigalow Belt South Bioregions	Not detected	0
New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Escarpment Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Northern Escarpment Wet Sclerophyll Forests greater than or equal to 70% and less than 90%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Low	<1
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Western Slopes Grassy Woodlands greater than or equal to 70% and less than 90%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt	Moderate	1,305 (641 to 1,969)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions		

Table 9 New England Tablelands Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Armidale Plateau		
Clarence Dry Sclerophyll Forests less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Montane Bogs and Fens greater than or equal to 70% and less than 90%	Low	<1
New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	316 (198 to 433)
New England Dry Sclerophyll Forests less than 50%	Low	82 (52 to 113)
New England Grassy Woodlands greater than or equal to 50% and less than 70%	Low	24 (10 to 37)
New England Grassy Woodlands greater than or equal to 70% and less than 90%	Low	976 (427 to 1,525)
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	32 (17 to 46)
Northern Tableland Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Low	101 (63 to 139)
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	9 (6 to 12)
Northern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Low	520 (285 to 755)
Temperate Montane Grasslands greater than or equal to 50% and less than 70%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	561 (246 to 877)
Bundarra Downs		
New England Grassy Woodlands less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Deepwater Downs		
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Howell Shrublands in the New England Tableland and Nandewar Bioregions	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Eastern Nandewars		
Eastern Riverine Forests less than 50%	Not detected	0
Howell Shrublands in the New England Tableland and Nandewar Bioregions	Low	494 (308 to 680)
New England Grassy Woodlands greater than or equal to 50% and less than 70%	Low	2 (1 to 2)
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Montane Heaths less than 50%	Not detected	0
Northern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Tableland Clay Grassy Woodlands greater than or equal to 70% and less than 90%	Low	53 (29 to 77)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	310 (152 to 468)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Glenn Innes-Guyra Basalts		
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Low	114 (62 to 166)
Eastern Riverine Forests less than 50%	Not detected	0
New England Dry Sclerophyll Forests less than 50%	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	2 (1 to 3)
Northern Tableland Dry Sclerophyll Forests less than 50%	Low	3 (2 to 4)
Northern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Low	886 (485 to 1,286)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina	Moderate	1,148 (561 to 1,735)
Nightcap		
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Low	3 (2 to 5)
Round Mountain		
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	30 (16 to 44)
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Low	75 (41 to 108)
Severn River Volcanics		
Eastern Riverine Forests less than 50%	Not detected	0
Howell Shrublands in the New England Tableland and Nandewar Bioregions	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Montane Heaths less than 50%	Not detected	0
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions		
Stanthorpe Plateau		
New England Dry Sclerophyll Forests less than 50%	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
Tenterfield Plateau		
Eastern Riverine Forests less than 50%	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Walcha Plateau		
Clarence Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Mount Kaputar high elevation and dry rainforest land snail and slug community in the Nandewar and Brigalow Belt South Bioregions	Not detected	0
New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
New England Dry Sclerophyll Forests less than 50%	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Escarpment Wet Sclerophyll Forests greater than or equal to 70% and less than 90%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Northern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Yetman Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Wongwibinda Plateau		
New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	596 (375 to 817)
New England Dry Sclerophyll Forests less than 50%	Low	61 (38 to 83)
New England Grassy Woodlands greater than or equal to 70% and less than 90%	Low	55 (24 to 85)
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Low	229 (126 to 333)
Northern Gorge Dry Sclerophyll Forests less than 50%	Low	5 (2 to 8)
Northern Montane Heaths less than 50%	Low	11 (10 to 12)
Northern Tableland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	910 (584 to 1,236)
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Low	191 (105 to 278)
Yarrowyck-Kentucky Downs		
Howell Shrublands in the New England Tableland and Nandewar Bioregions	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion	Not detected	0
Northern Tableland Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Low	120 (75 to 165)
Northern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Northern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Ribbon Gum—Mountain Gum—Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion	Low	9 (5 to 13)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0

Table 10 NSW North Coast Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Carrai Plateau		
Carex Sedgeland of the New England Tableland, Nandewar, Brigalow Belt South and NSW North Coast Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Northern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	358 (189 to 527)
Northern Montane Heaths less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests less than 50%	Low	894 (573 to 1,214)
Coffs Coast and Escarpment		
Clarence Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	67 (37 to 97)
Coastal Heath Swamps less than 50%	Not detected	0
Coastal Swamp Forests less than 50%	Not detected	0
Littoral Rainforest in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	57 (31 to 83)
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Low	17 (6 to 28)
Maritime Grasslands greater than or equal to 70% and less than 90%	Low	51 (27 to 75)
North Coast Wet Sclerophyll Forests less than 50%	Low	75 (40 to 111)
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	5 (3 to 7)
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	7 (4 to 10)
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Ellerston		
Central Hunter Grey Box—Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions	Not detected	0
Central Hunter Ironbark—Spotted Gum—Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions	Low	4 (2 to 7)
Coastal Floodplain Wetlands greater than or equal to 70% and less than 90%	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Hunter Valley Vine Thicket in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	19 (11 to 27)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Not detected	0
Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions	Low	<1
Northern Hinterland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	47 (25 to 69)
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Low	5 (3 to 7)
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Western Slopes Grassy Woodlands greater than or equal to 50% and less than 70%	Low	<1
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	9 (4 to 13)
Karuah Manning		
Coastal Dune Dry Sclerophyll Forests less than 50%	Low	604 (358 to 849)
Coastal Heath Swamps less than 50%	Not detected	0
Coastal Swamp Forests less than 50%	Low	<1
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Dry Rainforests greater than or equal to 70% and less than 90%	Low	11 (5 to 17)
Dry Rainforests less than 50%	Low	22 (10 to 34)
Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions	Not detected	0
Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	56 (32 to 79)
Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Low	7 (4 to 10)
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Low	128 (75 to 181)
Kincumber Scribbly Gum Forest in the Sydney Basin Bioregion	Not detected	0
Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions	Low	61 (36 to 87)
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Low	52 (24 to 80)
Mangrove Swamps greater than or equal to 50% and less than 70%	Low	6 (4 to 8)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Maritime Grasslands greater than or equal to 70% and less than 90%	Not detected	0
North Coast Wet Sclerophyll Forests less than 50%	Low	6 (3 to 9)
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	165 (87 to 243)
South Coast Sands Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	<1
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	9 (6 to 13)
Sydney Freshwater Wetlands in the Sydney Basin Bioregion	Not detected	0
Umina Coastal Sandplain Woodland in the Sydney Basin Bioregion	Not detected	0
Wallum Sand Heaths less than 50%	Not detected	0
Macleay Gorges		
Clarence Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Strong	7,394 (4,075 to 10,711)
Coastal Valley Grassy Woodlands greater than or equal to 70% and less than 90%	Low	268 (141 to 395)
Eastern Riverine Forests less than 50%	Low	100 (54 to 146)
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Low	52 (24 to 80)
Northern Gorge Dry Sclerophyll Forests less than 50%	Moderate	3,603 (1,300 to 5,905)
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	208 (110 to 307)
Northern Montane Heaths less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests less than 50%	Moderate	3,259 (2,091 to 4,427)
Macleay Hastings		
Clarence Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	16 (9 to 23)
Clarence Dry Sclerophyll Forests less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Low	6 (3 to 9)
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Mangrove Swamps less than 50%	Not detected	0
North Coast Wet Sclerophyll Forests less than 50%	Low	<1
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Tomalia		
Central Hunter Grey Box—Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions	Not detected	0
Central Hunter Ironbark—Spotted Gum—Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions	Not detected	0
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Dry Rainforests less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Hunter Valley Vine Thicket in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Not detected	0
Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions	Not detected	0
Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
New England Dry Sclerophyll Forests less than 50%	Not detected	0
New England Grassy Woodlands less than 50%	Not detected	0
Northern Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Upper Hunter		
Central Hunter Ironbark—Spotted Gum—Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions	Not detected	0
Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions	Low	22 (13 to 31)
Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	347 (202 to 491)
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Low	559 (326 to 792)
Kurri Sand Swamp Woodland in the Sydney Basin Bioregion	Not detected	0
Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions	Low	877 (511 to 1,242)
Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
North Coast Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	13 (7 to 20)
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Low	19 (10 to 28)
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Yuraygir		
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0

Table 11 NSW South Western Slopes Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Capertee Valley		

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Western Slopes Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Slopes		
Alpine Herbfields less than 50%	Moderate	2,082 (1,143 to 3,020)
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests greater than or equal to 70% and less than 90%	Low	98 (53 to 142)
Eastern Riverine Forests less than 50%	Low	12 (6 to 17)
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Low	3 (2 to 5)
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Low	338 (191 to 484)
Inland Riverine Forests greater than or equal to 50% and less than 70%	Moderate	1,382 (783 to 1,980)
Inland Riverine Forests less than 50%	Moderate	3,633 (2,059 to 5,206)
Inland Rocky Hill Woodlands less than 50%	Moderate	1,216 (761 to 1,670)
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Low	<1
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	<1
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	297 (176 to 419)
Southern Tableland Dry Sclerophyll Forests less than 50%	Strong	5,550 (3,279 to 7,822)
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Low	<1
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	6 (3 to 8)
Southern Tableland Grassy Woodlands greater than or equal to 90%	Low	2 (1 to 4)
Southern Tableland Wet Sclerophyll Forests less than 50%	Low	4 (2 to 7)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Not detected	0
Temperate Montane Grasslands less than 50%	Low	45 (28 to 61)
Upper Riverina Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	174 (108 to 239)
Upper Riverina Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Not detected	0
Upper Riverina Dry Sclerophyll Forests greater than or equal to 90%	Moderate	2,460 (1,533 to 3,386)
Upper Riverina Dry Sclerophyll Forests less than 50%	Moderate	3,058 (1,906 to 4,209)
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Moderate	3,854 (2,653 to 5,054)
Western Slopes Dry Sclerophyll Forests greater than or equal to 90%	Low	8 (6 to 11)
Western Slopes Dry Sclerophyll Forests less than 50%	Low	768 (529 to 1,007)
Western Slopes Grassy Woodlands greater than or equal to 50% and less than 70%	Moderate	4,223 (2,065 to 6,383)
Western Slopes Grassy Woodlands less than 50%	Moderate	2,246 (1,098 to 3,394)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Strong	45,632 (22,548 to 68,725)
Lower Slopes		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Low	53 (30 to 76)
Floodplain Transition Woodlands greater than or equal to 70% and less than 90%	Low	3 (2 to 4)
Inland Floodplain Shrublands less than 50%	Low	31 (20 to 42)
Inland Floodplain Swamps greater than or equal to 50% and less than 70%	Strong	7,593 (3,622 to 11,563)
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Moderate	1,927 (1,091 to 2,763)
Inland Riverine Forests greater than or equal to 50% and less than 70%	Low	310 (176 to 444)
Inland Riverine Forests less than 50%	Not detected	0
Inland Rocky Hill Woodlands less than 50%	Low	44 (28 to 61)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Mallee and Mallee-Broombush dominated woodland and shrubland, lacking Triodia, in the NSW South Western Slopes Bioregion	Not detected	0
Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes Bioregions	Low	227 (144 to 310)
Riverine Plain Grasslands greater than or equal to 50% and less than 70%	Low	353 (290 to 416)
Western Peneplain Woodlands less than 50%	Not detected	0
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	36 (25 to 48)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0

Table 12 Riverina Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Lachlan		
Acacia loderi Shrublands	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Inland Floodplain Shrublands greater than or equal to 50% and less than 70%	Not detected	0
Inland Floodplain Swamps less than 50%	Not detected	0
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Saline Lakes less than 50%	Not detected	0
Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes Bioregions	Not detected	0
Riverine Chenopod Shrublands greater than or equal to 50% and less than 70%	Not detected	0
Riverine Plain Grasslands less than 50%	Not detected	0
Western Peneplain Woodlands less than 50%	Not detected	0
Murray Fans		
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Inland Floodplain Shrublands greater than or equal to 50% and less than 70%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Inland Floodplain Swamps greater than or equal to 50% and less than 70%	Not detected	0
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Low	28 (17 to 39)
Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions	Low	13 (8 to 19)
Riverine Chenopod Shrublands less than 50%	Not detected	0
Riverine Plain Grasslands greater than or equal to 70% and less than 90%	Low	22 (18 to 26)
Sandhill Pine Woodland in the Riverina, Murray-Darling Depression and NSW South Western Slopes Bioregions	Not detected	0
Murray Scroll Belt		
Acacia loderi Shrublands	Not detected	0
Aeolian Chenopod Shrublands less than 50%	Not detected	0
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Saline Lakes less than 50%	Not detected	0
Murrumbidgee		
Acacia melvillei Shrubland in the Riverina and Murray-Darling Depression Bioregions	Not detected	0
Aeolian Chenopod Shrublands less than 50%	Low	602 (387 to 816)
Floodplain Transition Woodlands greater than or equal to 50% and less than 70%	Moderate	2,285 (1,260 to 3,309)
Inland Floodplain Shrublands greater than or equal to 50% and less than 70%	Moderate	3,865 (2,510 to 5,222)
Inland Floodplain Shrublands less than 50%	Moderate	1,814 (1,188 to 2,441)
Inland Floodplain Swamps greater than or equal to 50% and less than 70%	Moderate	2,881 (1,374 to 4,387)
Inland Floodplain Swamps less than 50%	Low	87 (41 to 132)
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Moderate	3,048 (1,859 to 4,237)
Inland Riverine Forests greater than or equal to 50% and less than 70%	Low	28 (16 to 41)
Inland Riverine Forests less than 50%	Low	282 (160 to 404)
Inland Saline Lakes less than 50%	Moderate	1,198 (745 to 1,653)
Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes Bioregions	Strong	35,386 (27,707 to 43,066)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Riverine Chenopod Shrublands greater than or equal to 50% and less than 70%	Strong	45,626 (40,214 to 51,033)
Riverine Chenopod Shrublands less than 50%	Strong	200,933 (177,110 to 224,733)
Riverine Plain Grasslands greater than or equal to 50% and less than 70%	Strong	5,720 (4,697 to 6,742)
Riverine Plain Grasslands greater than or equal to 70% and less than 90%	Strong	124,696 (102,409 to 146,984)
Riverine Plain Grasslands less than 50%	Strong	63,163 (51,874 to 74,452)
Sandhill Pine Woodland in the Riverina, Murray-Darling Depression and NSW South Western Slopes Bioregions	Moderate	1,293 (717 to 1,869)
Robinvale Plains		
Acacia loderi Shrublands	Not detected	0
Aeolian Chenopod Shrublands less than 50%	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Dune Mallee Woodlands less than 50%	Not detected	0
Inland Floodplain Swamps less than 50%	Not detected	0
Inland Floodplain Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Inland Riverine Forests less than 50%	Not detected	0
Inland Saline Lakes less than 50%	Not detected	0
Tecticornia lylei, Wiry Glasswort, low open-shrubland in the Murray Darling Depression Bioregion	Not detected	0

Table 13 Simpson Strzelecki Dunefields Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Strzelecki Desert		
Acacia loderi Shrublands	Not detected	0
Aeolian Chenopod Shrublands less than 50%	Not detected	0
Artesian Springs Ecological Community in the Great Artesian Basin	Not detected	0
Gibber Chenopod Shrublands less than 50%	Not detected	0
Gibber Transition Shrublands less than 50%	Not detected	0
Inland Floodplain Shrublands greater than or equal to 50% and less than 70%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Inland Floodplain Shrublands less than 50%	Not detected	0
Inland Saline Lakes less than 50%	Not detected	0
North-west Floodplain Woodlands less than 50%	Not detected	0
Sand Plain Mulga Shrublands greater than or equal to 50% and less than 70%	Not detected	0
Sand Plain Mulga Shrublands less than 50%	Not detected	0
Stony Desert Mulga Shrublands less than 50%	Not detected	0

Table 14 South East Corner Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Bateman		
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Illawarra Lowlands Grassy Woodland in the Sydney Basin Bioregion	Not detected	0
South Coast Wet Sclerophyll Forests less than 50%	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Lowland Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Warm Temperate Rainforests less than 50%	Not detected	0
South East Coastal Ranges		
Bangalay Sand Forest of the Sydney Basin and South East Corner Bioregions	Not detected	0
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Mangrove Swamps less than 50%	Not detected	0
South Coast Heaths less than 50%	Not detected	0
South Coast Wet Sclerophyll Forests less than 50%	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Lowland Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Warm Temperate Rainforests less than 50%	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0

Table 15 South Eastern Highlands Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Bathurst		
Central Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Low	<1
Eastern Riverine Forests less than 50%	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	151 (89 to 213)
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Grassy Woodlands less than 50%	Not detected	0
Sydney Montane Dry Sclerophyll Forests less than 50%	Not detected	0
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Moderate	1,322 (723 to 1,921)
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Tableland Clay Grassy Woodlands less than 50%	Not detected	0
Upper Riverina Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	47 (30 to 65)
Upper Riverina Dry Sclerophyll Forests less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	565 (284 to 846)
Bondo		
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Low	411 (224 to 597)
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	128 (70 to 186)
Southern Tableland Wet Sclerophyll Forests less than 50%	Low	158 (83 to 232)
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Not detected	0
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Low	8 (4 to 11)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Upper Riverina Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	41 (22 to 60)
Bungonia		
Central Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps Bioregions	Low	2 (1 to 3)
Mt Canobolas Xanthoparmelia Lichen Community	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Low	208 (138 to 279)
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	487 (287 to 686)
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 90%	Low	2 (1 to 3)
Southern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
Temperate Montane Grasslands greater than or equal to 50% and less than 70%	Not detected	0
Werriwa Tablelands Cool Temperate Grassy Woodland in the South Eastern Highlands and South East Corner Bioregions	Low	116 (73 to 159)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	22 (12 to 31)
Capertee Uplands		
Central Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Corner, South Eastern Highlands and Australian Alps Bioregions		
Mt Canobolas Xanthoparmelia Lichen Community	Not detected	0
Newnes Plateau Shrub Swamp in the Sydney Basin Bioregion	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Grassy Woodlands less than 50%	Not detected	0
Subalpine Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Sydney Montane Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Montane Heaths less than 50%	Not detected	0
Tableland Clay Grassy Woodlands less than 50%	Not detected	0
Upper Riverina Dry Sclerophyll Forests less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Crookwell		
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Southern Escarpment Wet Sclerophyll Forests less than 50%	Low	<1
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	77 (46 to 109)
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Low	4 (2 to 6)
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	11 (6 to 16)
Southern Tableland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	2 (1 to 4)
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Low	594 (313 to 875)
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Tableland Clay Grassy Woodlands less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Upper Riverina Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	4 (2 to 5)
Werriwa Tablelands Cool Temperate Grassy Woodland in the South Eastern Highlands and South East Corner Bioregions	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Hill End		
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Low	364 (198 to 530)
Eastern Riverine Forests less than 50%	Not detected	0
Mt Canobolas Xanthoparmelia Lichen Community	Not detected	0
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Strong	5,797 (3,424 to 8,169)
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Low	543 (297 to 790)
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	499 (273 to 725)
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Low	314 (172 to 457)
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Low	16 (9 to 23)
Upper Riverina Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Moderate	3,071 (1,915 to 4,228)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	528 (287 to 769)
Monaro		
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Monaro Tableland Cool Temperate Grassy Woodland in the South Eastern Highlands Bioregion	Low	36 (20 to 52)
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps Bioregions	Not detected	0
Mt Canobolas Xanthoparmelia Lichen Community	Not detected	0
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Low	580 (317 to 842)
Southern Tableland Grassy Woodlands less than 50%	Not detected	0
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Low	12 (6 to 17)
Tableland Clay Grassy Woodlands less than 50%	Not detected	0
Temperate Montane Grasslands greater than or equal to 70% and less than 90%	Strong	14,344 (9,030 to 19,661)
Upland Wetlands of the Drainage Divide of the New England Tableland Bioregion	Not detected	0
Upper Riverina Dry Sclerophyll Forests less than 50%	Not detected	0
Werriwa Tablelands Cool Temperate Grassy Woodland in the South Eastern Highlands and South East Corner Bioregions	Low	28 (18 to 38)
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Murrumbateman		
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Low	2 (1 to 3)
Eastern Riverine Forests less than 50%	Not detected	0
Inland Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Mt Canobolas Xanthoparmelia Lichen Community	Not detected	0
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	116 (63 to 169)
Southern Tableland Grassy Woodlands less than 50%	Not detected	0
Tableland Clay Grassy Woodlands less than 50%	Not detected	0
Upper Riverina Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Upper Riverina Dry Sclerophyll Forests greater than or equal to 70% and less than 90%	Low	3 (2 to 4)
Upper Riverina Dry Sclerophyll Forests less than 50%	Not detected	0
Werriwa Tablelands Cool Temperate Grassy Woodland in the South Eastern Highlands and South East Corner Bioregions	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North	Low	114 (62 to 165)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions		
Oberon		
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	<1
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	1 (1 to 2)
Subalpine Woodlands greater than or equal to 70% and less than 90%	Low	12 (7 to 16)
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Low	376 (202 to 550)
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Low	23 (13 to 34)
Tableland Clay Grassy Woodlands less than 50%	Not detected	0
Orange		
Carex Sedgeland of the New England Tableland, Nandewar, Brigalow Belt South and NSW North Coast Bioregions	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Monaro Tableland Cool Temperate Grassy Woodland in the South Eastern Highlands Bioregion	Not detected	0
Mt Canobolas Xanthoparmelia Lichen Community	Not detected	0
Southern Montane Heaths greater than or equal to 50% and less than 70%	Low	14 (8 to 19)
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 70% and less than 90%	Low	48 (26 to 70)
Southern Tableland Grassy Woodlands greater than or equal to 90%	Low	<1
Southern Tableland Grassy Woodlands less than 50%	Not detected	0
Southern Tableland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	21 (11 to 32)
Subalpine Woodlands less than 50%	Low	324 (201 to 447)
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Not detected	0
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Low	<1
Tableland Clay Grassy Woodlands less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Upper Riverina Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	2 (1 to 3)
Upper Riverina Dry Sclerophyll Forests less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0

Table 16 South Eastern Queensland Bioregion: forecast longer-term demand (at 28 March 2023)

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Burringbar-Conondale Ranges		
Coastal Cypress Pine Forest in the New South Wales North Coast Bioregion	Not detected	0
Coastal Dune Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Freshwater Lagoons less than 50%	Not detected	0
Coastal Headland Heaths less than 50%	Not detected	0
Coastal Heath Swamps less than 50%	Not detected	0
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Coastal Swamp Forests greater than or equal to 70% and less than 90%	Not detected	0
Coastal Swamp Forests less than 50%	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Littoral Rainforest in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Lowland Rainforest on Floodplain in the New South Wales North Coast Bioregion	Low	<1
Mangrove Swamps greater than or equal to 50% and less than 70%	Low	7 (4 to 9)
North Coast Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
North Coast Wet Sclerophyll Forests less than 50%	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	11 (7 to 16)
Sydney Freshwater Wetlands in the Sydney Basin Bioregion	Not detected	0
Temperate Montane Grasslands greater than or equal to 50% and less than 70%	Not detected	0
Wallum Sand Heaths less than 50%	Low	2 (1 to 3)
Clarence Lowlands		
Clarence Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	317 (174 to 459)
Clarence Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Freshwater Lagoons less than 50%	Not detected	0
Coastal Headland Heaths less than 50%	Not detected	0
Coastal Heath Swamps less than 50%	Not detected	0
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Coastal Swamp Forests greater than or equal to 70% and less than 90%	Low	<1
Coastal Swamp Forests less than 50%	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	<1
Littoral Rainforest in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Low	8 (3 to 13)
Maritime Grasslands greater than or equal to 70% and less than 90%	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Low	719 (396 to 1,042)
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	60 (37 to 84)
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	89 (54 to 125)
Sydney Freshwater Wetlands in the Sydney Basin Bioregion	Not detected	0
Themeda grassland on seacliffs and coastal headlands in the NSW North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Wallum Sand Heaths less than 50%	Not detected	0
Clarence Sandstones		
North Coast Dry Sclerophyll Forests less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0
Scenic Rim		
Coastal Valley Grassy Woodlands less than 50%	Low	3 (2 to 5)
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Low	227 (117 to 338)
North Coast Dry Sclerophyll Forests less than 50%	Low	41 (19 to 63)
North Coast Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	4 (2 to 5)
North Coast Wet Sclerophyll Forests less than 50%	Moderate	1,858 (988 to 2,727)
Northern Hinterland Wet Sclerophyll Forests less than 50%	Low	663 (350 to 975)
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	<1
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0

Table 17 Sydney Basin Bioregion: forecast longer-term demand (sat 28 March 2023)

	<u>-</u>	
IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Burragorang		
Blue Mountains Shale Cap Forest in the Sydney Basin Bioregion	Not detected	0
Central Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Floodplain Wetlands less than 50%	Not detected	0
Cumberland Plain Woodland in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Northern Warm Temperate Rainforests less than 50%	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Lowland Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Hinterland Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Montane Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Sand Flats Dry Sclerophyll Forests less than 50%	Not detected	0
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Cumberland		
Blue Mountains Shale Cap Forest in the Sydney Basin Bioregion	Not detected	0
Castlereagh Scribbly Gum Woodland in the Sydney Basin Bioregion	Low	202 (133 to 271)
Castlereagh Swamp Woodland Community	Low	<1
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Cooks River/Castlereagh Ironbark Forest in the Sydney Basin Bioregion	Low	21 (11 to 30)
Cumberland Plain Woodland in the Sydney Basin Bioregion	Moderate	2,782 (1,467 to 4,097)
Dry Rainforests less than 50%	Not detected	0
Duffys Forest Ecological Community in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Hygrocybeae Community of Lane Cove Bushland Park in the Sydney Basin Bioregion	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Mangrove Swamps less than 50%	Not detected	0
Moist Shale Woodland in the Sydney Basin Bioregion	Not detected	0
Northern Hinterland Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	4 (2 to 6)
Northern Warm Temperate Rainforests greater than or equal to 70% and less than 90%	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Moderate	1,926 (1,056 to 2,796)
Shale Gravel Transition Forest in the Sydney Basin Bioregion	Not detected	0
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	4 (2 to 6)
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	<1
Sydney Coastal Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	1 (1 to 1)
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Coastal Heaths less than 50%	Not detected	0
Sydney Hinterland Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion	Low	3 (2 to 5)
Tableland Clay Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Ettrema		
Blue Mountains Shale Cap Forest in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Not detected	0
Illawarra Subtropical Rainforest in the Sydney Basin Bioregion	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Lowland Wet Sclerophyll Forests less than 50%	Not detected	0
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Hunter		

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Central Hunter Grey Box—Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions	Moderate	2,813 (1,640 to 3,987)
Central Hunter Ironbark—Spotted Gum—Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions	Strong	10,428 (5,499 to 15,358)
Coastal Dune Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Floodplain Wetlands greater than or equal to 70% and less than 90%	Low	187 (103 to 272)
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	40 (22 to 57)
Coastal Valley Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Coastal Valley Grassy Woodlands greater than or equal to 70% and less than 90%	Low	<1
Dry Rainforests less than 50%	Not detected	0
Eastern Riverine Forests greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests greater than or equal to 70% and less than 90%	Low	40 (22 to 58)
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Not detected	0
Hunter Floodplain Red Gum Woodland in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions	Low	306 (161 to 450)
Hunter Valley Footslopes Slaty Gum Woodland in the Sydney Basin Bioregion	Not detected	0
Hunter Valley Vine Thicket in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Hunter Valley Weeping Myall Woodland in the Sydney Basin Bioregion	Not detected	0
Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	<1
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Not detected	0
Kincumber Scribbly Gum Forest in the Sydney Basin Bioregion	Not detected	0
Kurri Sand Swamp Woodland in the Sydney Basin Bioregion	Not detected	0
Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions	Moderate	1,141 (665 to 1,617)
Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Mangrove Swamps greater than or equal to 50% and less than 70%	Low	287 (187 to 388)
Mangrove Swamps less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Low	232 (134 to 330)
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Not Classified	Not detected	<1
Pittwater and Wagstaffe Spotted Gum Forest in the Sydney Basin Bioregion	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	968 (531 to 1,404)
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Coastal Heaths less than 50%	Not detected	0
Umina Coastal Sandplain Woodland in the Sydney Basin Bioregion	Not detected	0
Upland Wetlands of the Drainage Divide of the New England Tableland Bioregion	Not detected	0
Warkworth Sands Woodland in the Sydney Basin Bioregion	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Low	211 (103 to 318)
Illawarra		
Bangalay Sand Forest of the Sydney Basin and South East Corner Bioregions	Low	47 (28 to 66)
Coastal Floodplain Wetlands greater than or equal to 90%	Not detected	0
Coastal Headland Heaths greater than or equal to 50% and less than 70%	Low	2 (1 to 4)
Coastal Headland Heaths less than 50%	Not detected	0
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Coastal Upland Swamp in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Illawarra Lowlands Grassy Woodland in the Sydney Basin Bioregion	Not detected	0
Illawarra Subtropical Rainforest in the Sydney Basin Bioregion	Low	27 (12 to 41)
Maritime Grasslands greater than or equal to 70% and less than 90%	Not detected	0
Melaleuca armillaris Tall Shrubland in the Sydney Basin Bioregion	Low	82 (50 to 113)
North Coast Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Warm Temperate Rainforests less than 50%	Not detected	0
South Coast Wet Sclerophyll Forests less than 50%	Not detected	0
South East Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Lowland Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Warm Temperate Rainforests less than 50%	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Coastal Heaths less than 50%	Not detected	0
Sydney Hinterland Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Montane Dry Sclerophyll Forests less than 50%	Not detected	0
Themeda grassland on seacliffs and coastal headlands in the NSW North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Jervis		
Illawarra Lowlands Grassy Woodland in the Sydney Basin Bioregion	Not detected	0
Illawarra Subtropical Rainforest in the Sydney Basin Bioregion	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
South Coast Wet Sclerophyll Forests less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Kerrabee		
Central Gorge Dry Sclerophyll Forests less than 50%	Low	2 (1 to 4)
Central Hunter Grey Box—Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions	Low	639 (371 to 907)
Coastal Floodplain Wetlands greater than or equal to 70% and less than 90%	Not detected	0
Coastal Valley Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Fuzzy Box Woodland on alluvial Soils of the South Western Slopes, Darling Riverine Plains and Brigalow Belt South Bioregions	Not detected	0
Hunter Floodplain Red Gum Woodland in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Hunter Valley Weeping Myall Woodland in the Sydney Basin Bioregion	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70%	Low	119 (69 to 170)
North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 70% and less than 90%	Low	7 (4 to 11)
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	213 (146 to 279)
Western Slopes Dry Sclerophyll Forests less than 50%	Low	318 (219 to 417)
Western Slopes Grassy Woodlands greater than or equal to 90%	Low	165 (81 to 249)
Western Slopes Grassy Woodlands less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Moderate	2,969 (1,452 to 4,487)
Moss Vale		
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Southern Highlands Shale Woodlands in the Sydney Basin Bioregion	Not detected	0
Southern Tableland Wet Sclerophyll Forests less than 50%	Not detected	0
Pittwater		
Bangalay Sand Forest of the Sydney Basin and South East Corner Bioregions	Not detected	0
Blue Gum High Forest in the Sydney Basin Bioregion	Not detected	0
Coastal Dune Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Headland Heaths less than 50%	Not detected	0
Coastal Saltmarsh in the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Coastal Upland Swamp in the Sydney Basin Bioregion	Not detected	0
Cooks River/Castlereagh Ironbark Forest in the Sydney Basin Bioregion	Not detected	0
Duffys Forest Ecological Community in the Sydney Basin Bioregion	Low	4 (2 to 6)
Eastern Suburbs Banksia Scrub in the Sydney Basin Bioregion	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Hygrocybeae Community of Lane Cove Bushland Park in the Sydney Basin Bioregion	Not detected	0
Kurnell Dune Forest in the Sutherland Shire and City of Rockdale	Not detected	0
Mangrove Swamps less than 50%	Not detected	0
Maritime Grasslands greater than or equal to 70% and less than 90%	Not detected	0
North Coast Wet Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Northern Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests greater than or equal to 90%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Warm Temperate Rainforests less than 50%	Not detected	0
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Sydney Coastal Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Sydney Coastal Dry Sclerophyll Forests greater than or equal to 90%	Not detected	0
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Coastal Heaths greater than or equal to 70% and less than 90%	Not detected	0
Sydney Coastal Heaths less than 50%	Not detected	0
Sydney Hinterland Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion	Low	2 (1 to 3)
Sydney Cataract		
Bangalay Sand Forest of the Sydney Basin and South East Corner Bioregions	Not detected	0
Blue Mountains Shale Cap Forest in the Sydney Basin Bioregion	Not detected	0
Coastal Upland Swamp in the Sydney Basin Bioregion	Not detected	0
Cumberland Plain Woodland in the Sydney Basin Bioregion	Not detected	0
Dry Rainforests less than 50%	Not detected	0
Duffys Forest Ecological Community in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Hygrocybeae Community of Lane Cove Bushland Park in the Sydney Basin Bioregion	Not detected	0
Illawarra Lowlands Grassy Woodland in the Sydney Basin Bioregion	Not detected	0
Illawarra Subtropical Rainforest in the Sydney Basin Bioregion	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
North Coast Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Warm Temperate Rainforests less than 50%	Not detected	0
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Not detected	0
South Coast Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Lowland Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Sydney sheltered forest on transitional sandstone soils in the Sydney Basin Bioregion	Not detected	0
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Coastal Heaths less than 50%	Not detected	0
Sydney Hinterland Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Montane Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion	Not detected	0
Wollemi		
Blue Mountains Swamps in the Sydney Basin Bioregion	Low	<1
Central Gorge Dry Sclerophyll Forests less than 50%	Not detected	0
Cumberland Plain Woodland in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps Bioregions	Not detected	0
Newnes Plateau Shrub Swamp in the Sydney Basin Bioregion	Not detected	0
North-west Slopes Dry Sclerophyll Woodlands less than 50%	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Not detected	0
Southern Escarpment Wet Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Dry Sclerophyll Forests less than 50%	Not detected	0
Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Southern Tableland Grassy Woodlands less than 50%	Not detected	0
Subalpine Woodlands greater than or equal to 50% and less than 70%	Not detected	0
Sydney Hinterland Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Montane Dry Sclerophyll Forests less than 50%	Low	38 (18 to 58)
Sydney Montane Heaths less than 50%	Not detected	0
Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion	Not detected	0
Tableland Basalt Forest in the Sydney Basin and South Eastern Highlands Bioregions	Not detected	0
Tableland Clay Grassy Woodlands less than 50%	Not detected	0
Upper Riverina Dry Sclerophyll Forests less than 50%	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Western Slopes Dry Sclerophyll Forests less than 50%	Not detected	0
Western Slopes Grassy Woodlands less than 50%	Not detected	0
White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions	Not detected	0
Wyong		
Bangalay Sand Forest of the Sydney Basin and South East Corner Bioregions	Not detected	0
Coastal Dune Dry Sclerophyll Forests less than 50%	Not detected	0
Coastal Headland Heaths less than 50%	Not detected	0
Dry Rainforests less than 50%	Not detected	0
Duffys Forest Ecological Community in the Sydney Basin Bioregion	Not detected	0
Hunter Lowland Redgum Forest in the Sydney Basin and New South Wales North Coast Bioregions	Not detected	0
Hunter-Macleay Dry Sclerophyll Forests less than 50%	Low	14 (8 to 20)
Hygrocybeae Community of Lane Cove Bushland Park in the Sydney Basin Bioregion	Not detected	0
Kincumber Scribbly Gum Forest in the Sydney Basin Bioregion	Not detected	0
Kurnell Dune Forest in the Sutherland Shire and City of Rockdale	Not detected	0
Kurri Sand Swamp Woodland in the Sydney Basin Bioregion	Not detected	0
Low woodland with heathland on indurated sand at Norah Head	Not detected	0
Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Maritime Grasslands greater than or equal to 70% and less than 90%	Not detected	0
North Coast Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Pittwater and Wagstaffe Spotted Gum Forest in the Sydney Basin Bioregion	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Subtropical Coastal Floodplain Forest of the New South Wales North Coast Bioregion	Not detected	0

IBRA subregion and Offset Trading Group	Indicative level of credit demand	Estimated number of credits
Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Low	28 (17 to 39)
Sydney Coastal Dry Sclerophyll Forests greater than or equal to 50% and less than 70%	Low	623 (385 to 861)
Sydney Coastal Dry Sclerophyll Forests less than 50%	Not detected	0
Sydney Freshwater Wetlands in the Sydney Basin Bioregion	Not detected	0
Themeda grassland on seacliffs and coastal headlands in the NSW North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Wallum Sand Heaths less than 50%	Not detected	0
Yengo		
Coastal Valley Grassy Woodlands less than 50%	Not detected	0
Cumberland Plain Woodland in the Sydney Basin Bioregion	Low	25 (13 to 37)
Dry Rainforests less than 50%	Not detected	0
Duffys Forest Ecological Community in the Sydney Basin Bioregion	Not detected	0
Eastern Riverine Forests less than 50%	Not detected	0
Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions	Not detected	0
Maroota Sands Swamp Forest	Not detected	0
Northern Hinterland Wet Sclerophyll Forests less than 50%	Not detected	0
Northern Warm Temperate Rainforests greater than or equal to 70% and less than 90%	Not detected	0
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions	Not detected	0
Shale Sandstone Transition Forest in the Sydney Basin Bioregion	Not detected	0
Sydney Coastal Dry Sclerophyll Forests less than 50%		
	Not detected	0
Sydney Coastal Heaths less than 50%	Not detected Not detected	0

Environment and Heritage (EHG), Department of Planning and Environment Locked Bag 5022, Parramatta NSW 2124; Phone: 1300 361 967 (EHG enquiries); Email: info@environment.nsw.gov.au; Website: www.environment.nsw.gov.au ISBN 978-1-923132-29-0; EHG 2023/0358; November 2023.