

Ecosystem credit demand estimate

Offset Trading Group indicative nearer-term demand by IBRA region and subregion

The following ecosystem credit demand estimate has been produced using publicly available information from the Major Projects Planning Portal and data from the Biodiversity Assessment Method Calculator system. This is presented as demand per Offset Trading Group by IBRA (Interim Biogeographic Regionalisation for Australia) subregion. The estimate is based on information as of 28 March 2023.

Landholders should seek their own legal, financial and ecological advice before participating in the market. As the method to generate this estimate uses data from stages of the planning process before approval has been granted, the resulting credit predictions associated with these major projects can be uncertain.

The data are presented in a series of tables, one for each IBRA bioregion as follows (click on the bioregion of interest to jump to the table):

| Table 1 | Brigalow Belt South Bioregion | 2 |
|----------|------------------------------------|---|
| Table 2 | Darling Riverine Plains Bioregion | 3 |
| Table 3 | Nandewar Bioregion | 3 |
| Table 4 | New England Tablelands Bioregion | 4 |
| Table 5 | NSW North Coast Bioregion | 5 |
| Table 6 | NSW South Western Slopes Bioregion | 6 |
| Table 7 | Riverina Bioregion | 7 |
| Table 8 | South Eastern Highlands Bioregion | 8 |
| Table 9 | South Eastern Queensland Bioregion | 9 |
| Table 10 | South Eastern Highlands Bioregion | 9 |

A numerical credit estimate is provided. The following categories have also been used to depict demand:

Not detected Low (>0 to 999) Moderate (1,000 to 4,999) Strong (>5,000) The ecosystem credit demand estimate Excel spreadsheet containing the same data is also available for download.

| Table 1 | Brigalow Belt South Bioregion: forecast nearer-term demand |
|---------|--|
| | (at 28 March 2023) |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Liverpool Plains | | |
| Pilliga Outwash Dry Sclerophyll Forests less than 50% | Low | 119 |
| Liverpool Range | | |
| Eastern Riverine Forests less than 50% | Low | 69 |
| New England Dry Sclerophyll Forests less than 50% | Low | 147 |
| New England Grassy Woodlands greater than or equal to 50% and less than 70% | Strong | 7,153 |
| New England Grassy Woodlands less than 50% | Low | 317 |
| Western Slopes Grassy Woodlands greater than or equal to 90% | Moderate | 3,948 |
| Western Slopes Grassy Woodlands less than 50% | Low | 669 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Strong | 10,216 |
| Northern Outwash | | |
| Carbeen Open Forest Community in the Darling Riverine Plains and Brigalow Belt South Bioregions | Low | 35 |
| Inland Riverine Forests greater than or equal to 50% and less than 70% | Low | 237 |
| Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes bioregions | Low | 156 |
| Semi-arid Floodplain Grasslands greater than or equal to 70% and less than 90% | Low | 351 |
| Pilliga | | |
| Eastern Riverine Forests less than 50% | Low | 36 |
| New England Grassy Woodlands greater than or equal to 50% and less than 70% | Low | 65 |
| North Coast Dry Sclerophyll Forests less than 50% | Low | 230 |
| South Coast Sands Dry Sclerophyll Forests less than 50% | Low | 237 |
| Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 747 |
| Western Slopes Dry Sclerophyll Forests less than 50% | Low | 605 |
| Western Slopes Grassy Woodlands greater than or equal to 90% | Moderate | 1,759 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Western Slopes Grassy Woodlands less than 50% | Low | 21 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Strong | 7,677 |

Table 2Darling Riverine Plains Bioregion: forecast nearer-term demand
(at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|---|---|-----------------------------------|
| Castlereagh-Barwon | | |
| Inland Riverine Forests greater than or equal to 50% and less than 70% | Low | 52 |
| Semi-arid Floodplain Grasslands greater than or equal to 70% and less than 90% | Strong | 5,688 |
| Wilcannia Plains | | |
| Coolibah-Black Box Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain and Mulga Lands Bioregions | Low | 63 |
| Inland Riverine Forests greater than or equal to 50% and less than 70% | Low | 45 |
| Riverine Chenopod Shrublands greater than or equal to 70% and less than 90% | Low | 139 |

Table 3 Nandewar Bioregion: forecast nearer-term demand (at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Peel | | |
| Eastern Riverine Forests less than 50% | Moderate | 1,771 |
| New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Moderate | 2,428 |
| New England Grassy Woodlands greater than or equal to 50% and less than 70% | Low | 128 |
| New England Grassy Woodlands less than 50% | Low | 69 |
| North-west Slopes Dry Sclerophyll Woodlands less than 50% | Moderate | 1,236 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Moderate | 1,295 |

Table 4New England Tablelands Bioregion: forecast nearer-term demand
(at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Armidale Plateau | | |
| Eastern Riverine Forests less than 50% | Low | 7 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Moderate | 1,213 |
| Eastern Nandewar | | |
| Carex Sedgeland of the New England Tableland, Nandewar, Brigalow Belt South and NSW North Coast Bioregions | Low | 71 |
| New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 143 |
| New England Dry Sclerophyll Forests greater than or equal to 70% and less than 90% | Low | 640 |
| New England Grassy Woodlands greater than or equal to 70% and less than 90% | Moderate | 3,239 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Low | 377 |
| Walcha Plateau | | |
| Montane Bogs and Fens greater than or equal to 70% and less than 90% | Low | 17 |
| New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Moderate | 2,243 |
| New England Dry Sclerophyll Forests less than 50% | Low | 386 |
| New England Grassy Woodlands greater than or equal to 50% and less than 70% | Moderate | 1,097 |
| New England Peppermint (Eucalyptus nova-anglica) Woodland on Basalts and Sediments in the New England Tableland Bioregion | Low | 396 |
| Northern Escarpment Wet Sclerophyll Forests greater than or equal to 70% and less than 90% | Low | 357 |
| Ribbon Gum–Mountain Gum–Snow Gum Grassy Forest/Woodland of the New England Tableland Bioregion | Low | 685 |
| Temperate Montane Grasslands greater than or equal to 70% and less than 90% | Low | 53 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney | Moderate | 3,962 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|---|---|-----------------------------------|
| Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | | |

Table 5 NSW North Coast Bioregion: forecast nearer-term demand (at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits | |
|--|---|-----------------------------------|--|
| Ellerston | | | |
| Central Hunter Ironbark–Spotted Gum–Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions | Low | 179 | |
| Dry Rainforests greater than or equal to 50% and less than 70% | Low | 26 | |
| Eastern Riverine Forests less than 50% | Low | 16 | |
| Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 175 | |
| Northern Hinterland Wet Sclerophyll Forests less than 50% | Low | 499 | |
| North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70% | Low | 10 | |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Moderate | 1,009 | |
| Karuah Manning | | | |
| Coastal Dune Dry Sclerophyll Forests less than 50% | Low | 439 | |
| Hunter-Macleay Dry Sclerophyll Forests less than 50% | Low | 325 | |
| Lower Hunter Spotted Gum Ironbark Forest in the Sydney Basin and NSW North Coast Bioregions | Low | 34 | |
| Lowland Rainforest in the NSW North Coast and Sydney Basin Bioregions | Low | 15 | |
| North Coast Wet Sclerophyll Forests less than 50% | Low | 231 | |
| Northern Hinterland Wet Sclerophyll Forests less than 50% | Low | 67 | |
| South Coast Sands Dry Sclerophyll Forests greater than or equal to 70% and less than 90% | Low | 37 | |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 353 | |
| Tomalla | | | |
| Eastern Riverine Forests less than 50% | Low | 77 | |
| Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 6 | |
| New England Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Moderate | 1,612 | |
| New England Grassy Woodlands less than 50% | Low | 106 | |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Northern Escarpment Wet Sclerophyll Forests greater than or equal to 70% and less than 90% | Low | 285 |
| Northern Gorge Dry Sclerophyll Forests less than 50% | Low | 157 |
| Northern Hinterland Wet Sclerophyll Forests less than 50% | Low | 288 |
| Northern Tableland Wet Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 54 |
| Northern Tableland Wet Sclerophyll Forests greater than or equal to 70% and less than 90% | Low | 691 |
| North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70% | Low | 27 |
| Tableland Clay Grassy Woodlands greater than or equal to 70% and less than 90% | Low | 3 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Moderate | 2,140 |
| Upper Hunter | | |
| Central Hunter Ironbark–Spotted Gum–Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions | Low | 3 |
| Eastern Riverine Forests less than 50% | Low | 4 |
| Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 10 |
| Lower Hunter Valley Dry Rainforest in the Sydney Basin and NSW North Coast Bioregions | Low | 21 |
| Northern Hinterland Wet Sclerophyll Forests less than 50% | Low | 38 |
| North-west Slopes Dry Sclerophyll Woodlands greater than or equal to 50% and less than 70% | Low | 1 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Low | 124 |

Table 6NSW South Western Slopes Bioregion: forecast nearer-term demand
(at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Inland Slopes | | |
| Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions | Moderate | 1,001 |
| Inland Riverine Forests less than 50% | Low | 190 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Inland Rocky Hill Woodlands greater than or equal to 50% and less than 70% | Low | 47 |
| Western Slopes Dry Sclerophyll Forests less than 50% | Low | 120 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Moderate | 2,960 |
| Lower Slopes | | |
| Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions | Low | 4 |
| Inland Riverine Forests greater than or equal to 50% and less than 70% | Low | 9 |
| Inland Riverine Forests less than 50% | Low | 1 |
| Western Slopes Grassy Woodlands greater than or equal to 90% | Low | 2 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Low | 333 |
| White Box Yellow Box Blakely's Red Gum Woodland | Low | 74 |

Table 7 Riverina Bioregion: forecast nearer-term demand (at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Murray Fans | | |
| Inland Grey Box Woodland in the Riverina, NSW South Western Slopes, Cobar Peneplain, Nandewar and Brigalow Belt South Bioregions | Low | 10 |
| Murrumbidgee | | |
| Inland Floodplain Shrublands greater than or equal to 50% and less than 70% | Low | 76 |
| Inland Floodplain Shrublands less than 50% | Low | 1 |
| Inland Floodplain Woodlands greater than or equal to 50% and less than 70% | Low | 30 |
| Inland Riverine Forests greater than or equal to 50% and less than 70% | Low | 3 |
| Inland Riverine Forests less than 50% | Low | 16 |
| Myall Woodland in the Darling Riverine Plains, Brigalow Belt South, Cobar Peneplain, Murray-Darling Depression, Riverina and NSW South Western Slopes Bioregions | Low | 797 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Riverine Plain Grasslands greater than or equal to 50% and less than 70% | Low | 80 |
| Riverine Plain Grasslands greater than or equal to 70% and less than 90% | Moderate | 1,773 |
| Riverine Plain Grasslands less than 50% | Moderate | 2,949 |
| Sandhill Pine Woodland in the Riverina, Murray-Darling Depression and NSW South Western Slopes Bioregions | Low | 145 |
| Robinvale Plains | | |
| Inland Floodplain Woodlands greater than or equal to 50% and less than 70% | Low | 18 |
| Sand Plain Mallee Woodlands less than 50% | Low | 65 |
| Sand Plain Mulga Shrublands less than 50% | Low | 19 |
| Semi-arid Sand Plain Woodlands greater than or equal to 50% and less than 70% | Low | 83 |

Table 8South Eastern Highlands Bioregion: forecast nearer-term demand
(at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Bathurst | | |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Low | 6 |
| Bungonia | | |
| Southern Tableland Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 7 |
| Temperate Montane Grasslands greater than or equal to 90% | Low | 26 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Low | 92 |
| Capertee Uplands | | |
| Southern Tableland Grassy Woodlands greater than or equal to 50% and less than 70% | Low | 29 |
| Subalpine Woodlands greater than or equal to 90% | Low | 8 |
| Monaro | | |
| Werriwa Tablelands Cool Temperate Grassy Woodland in the South Eastern Highlands and South East Corner Bioregions | Low | 31 |

Table 9South Eastern Queensland Bioregion: forecast nearer-term demand
(at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|---|---|-----------------------------------|
| Burringbar-Conondale Ranges | | |
| Coastal Freshwater Lagoons less than 50% | Low | 60 |
| Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 93 |

Table 10South Eastern Highlands Bioregion: forecast nearer-term demand
(at 28 March 2023)

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| Burragorang | | |
| Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps Bioregions | Low | 6 |
| Southern Escarpment Wet Sclerophyll Forests less than 50% | Low | 5 |
| Cumberland | | |
| Blue Gum High Forest in the Sydney Basin Bioregion | Low | 1 |
| Coastal Floodplain Wetlands greater than or equal to 50% and less than 70% | Low | 12 |
| Coastal Floodplain Wetlands greater than or equal to 90% | Low | 2 |
| Coastal Freshwater Lagoons greater than or equal to 70% and less than 90% | Low | 3 |
| Coastal Valley Grassy Woodlands greater than or equal to 90% | Low | 1 |
| Cumberland Dry Sclerophyll Forests greater than or equal to 90% | Low | 1 |
| Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest | Low | 36 |
| Cumberland Plain Woodland in the Sydney Basin Bioregion | Low | 194 |
| Eastern Riverine Forests less than 50% | Low | 2 |
| Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 4 |
| Mangrove Swamps greater than or equal to 70% and less than 90% | Low | 15 |
| River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 138 |
| Shale Gravel Transition Forest in the Sydney Basin Bioregion | Low | 40 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|---|---|-----------------------------------|
| Subtropical and Temperate Coastal Saltmarsh | Low | 2 |
| Swamp Oak Floodplain Forest of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 52 |
| Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion | Low | 11 |
| Ettrema | | |
| Eastern Riverine Forests greater than or equal to 50% and less than 70% | Low | 62 |
| Southern Lowland Wet Sclerophyll Forests less than 50% | Low | 39 |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 187 |
| Hunter | | |
| Central Hunter Grey Box–Ironbark Woodland in the New South Wales North Coast and Sydney Basin Bioregions | Moderate | 2,585 |
| Central Hunter Ironbark–Spotted Gum–Grey Box Forest in the New South Wales North Coast and Sydney Basin Bioregions | Low | 213 |
| Coastal Freshwater Lagoons greater than or equal to 70% and less than 90% | Low | 327 |
| Coastal Swamp Forests greater than or equal to 50% and less than 70% | Low | 74 |
| Coastal Valley Grassy Woodlands greater than or equal to 50% and less than 70% | Low | 672 |
| Coastal Valley Grassy Woodlands greater than or equal to 70% and less than 90% | Moderate | 1,297 |
| Eastern Riverine Forests greater than or equal to 50% and less than 70% | Low | 18 |
| Eastern Riverine Forests greater than or equal to 90% | Low | 13 |
| Eastern Riverine Forests less than 50% | Low | 3 |
| Freshwater Wetlands on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 4 |
| Hunter Floodplain Red Gum Woodland in the NSW North Coast and Sydney Basin Bioregions | Low | 41 |
| Hunter Valley Footslopes Slaty Gum Woodland in the Sydney Basin Bioregion | Low | 185 |
| Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 49 |
| Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 70% and less than 90% | Low | 201 |
| River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 2 |
| Warkworth Sands Woodland in the Sydney Basin Bioregion | Low | 59 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|--|---|-----------------------------------|
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Low | 253 |
| Illawarra | | |
| Illawarra Subtropical Rainforest in the Sydney Basin Bioregion | Low | 75 |
| Melaleuca armillaris Tall Shrubland in the Sydney Basin Bioregion | Low | 138 |
| North Coast Wet Sclerophyll Forests less than 50% | Low | 24 |
| South East Dry Sclerophyll Forests less than 50% | Low | 1 |
| Southern Lowland Wet Sclerophyll Forests less than 50% | Low | 8 |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 23 |
| Jervis | | |
| Coastal Heath Swamps less than 50% | Low | 39 |
| Kerrabee | | |
| Hunter Valley Footslopes Slaty Gum Woodland in the Sydney Basin Bioregion | Low | 342 |
| North Coast Dry Sclerophyll Forests less than 50% | Low | 437 |
| South Coast Sands Dry Sclerophyll Forests less than 50% | Low | 420 |
| Sydney Hinterland Dry Sclerophyll Forests less than 50% | Low | 9 |
| Western Slopes Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Moderate | 1,879 |
| Western Slopes Dry Sclerophyll Forests less than 50% | Moderate | 2,463 |
| Western Slopes Grassy Woodlands greater than or equal to 90% | Low | 9 |
| White Box–Yellow Box–Blakely's Red Gum Grassy Woodland and Derived Native Grassland in the NSW North Coast, New England Tableland, Nandewar, Brigalow Belt South, Sydney Basin, South Eastern Highlands, NSW South Western Slopes, South East Corner and Riverina Bioregions | Strong | 1,4587 |
| Moss Vale | | |
| Montane Bogs and Fens greater than or equal to 70% and less than 90% | Low | 6 |
| South East Dry Sclerophyll Forests less than 50% | Low | 148 |
| Southern Highlands Shale Woodlands in the Sydney Basin Bioregion | Low | 8 |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 96 |
| Pittwater | | |
| Duffys Forest Ecological Community in the Sydney Basin Bioregion | Low | 53 |

| IBRA subregion and Offset Trading Group | Indicative level of credit demand | Estimated number of credits |
|---|---|-----------------------------------|
| North Coast Wet Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 70 |
| Northern Hinterland Wet Sclerophyll Forests greater than or equal to 90% | Low | 19 |
| Sydney Coastal Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 2 |
| Sydney Coastal Dry Sclerophyll Forests greater than or equal to 90% | Low | 2 |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 128 |
| Sydney Coastal Heaths less than 50% | Low | 176 |
| Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion | Low | 1 |
| Wollemi | | |
| Sydney Montane Dry Sclerophyll Forests less than 50% | Low | 215 |
| Sydney Montane Heaths less than 50% | Low | 53 |
| Wyong | | |
| Coastal Freshwater Lagoons greater than or equal to 70% and less than 90% | Low | 5 |
| Hunter-Macleay Dry Sclerophyll Forests greater than or equal to 50% and less than 70% | Low | 3 |
| Swamp Sclerophyll Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions | Low | 10 |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 15 |
| Yengo | | |
| Sydney Coastal Dry Sclerophyll Forests less than 50% | Low | 248 |
| Sydney Hinterland Dry Sclerophyll Forests less than 50% | Low | 822 |

Environment and Heritage (EHG), Department of Planning and Environment Locked Bag 5022, Parramatta NSW 2124; Phone: 1300 361 967 (EHG enquiries); Email: info@environment.nsw.gov.au; Website: www.environment.nsw.gov.au ISBN 978-1-923132-32-0; EHG 2023/0362; November 2023.